



Confirmatory Composite Analysis

Florian Schuberth¹ Jörg Henseler¹
Theo K. Dijkstra²

¹University of Twente

²University of Groningen

June 18, 2019




Overview

- 1 Motivation
- 2 Confirmatory composite analysis
 - Model Specification
 - Model Identification
 - Model Estimation
 - Model Assessment
- 3 Monte Carlo simulation
- 4 Extensions

Behavioral concepts

Type of concept:	Behavioral concept
Type of construct:	Latent variable
Dominant statistical model:	Common factor model



Scientific paradigm:	Scientific Realism
Statistical approach:	Confirmatory factor analysis
Examples:	Abilities, attitudes, traits

Artifacts


Many disciplines deal with design concepts, so-called artifacts and their interplay with behavioral concepts:

Discipline	Behavioral concept	Artifact
Marketing:	Consumer brand attitude	Advertising mix
Criminology:	Intention to commit a crime	Prevention strategy
Education:	Pupil's knowledge base	Teaching program
Psychotherapy:	Mental illness	Psychiatric treatment

→ How to model and assess these artifacts?

Behavioral concepts & Artifacts

Type of concept:	Behavioral concept	Artifact
Type of construct:	Latent variable	Emergent variable
Dominant statistical model:	Common factor model	Composite model



Scientific paradigm:


Scientific Realism

Statistical approach:

Confirmatory factor analysis

Examples:

Abilities, attitudes, traits



Pragmatism

Confirmatory composite analysis


Indices, therapies,
intervention programs

Confirmatory Composite Analysis

Confirmatory composite analysis (CCA) consists of 4 steps:

- ① Specification of the composite model
- ② Identification of the composite model
- ③ Estimation of the composite model
- ④ Assessment of the composite model

Specification of the Composite Model



Minimal composite model

Is this a statistical model?

Consider the model-implied indicators' population covariance matrix:


$$\Sigma = \begin{pmatrix} \underline{x_1} & \underline{x_2} & \underline{y} & \underline{z} \\ \sigma_{11} & & & \\ \sigma_{12} & \sigma_{22} & & \\ \lambda_1 \sigma_{cy} & \lambda_2 \sigma_{cy} & \sigma_{yy} & \\ \lambda_1 \sigma_{cz} & \lambda_2 \sigma_{cz} & \sigma_{yz} & \sigma_{zz} \end{pmatrix},$$

where $\lambda_1 = \text{cov}(x_1, c)$ and $\lambda_2 = \text{cov}(x_2, c)$.


This matrix has rank-one constraints, which can be exploited in statistical testing.

→ Indeed, it is a statistical model

Composite Model vs. Common Factor Model



(a) Composite factor model



(b) Common factor model

Composite Model vs. Common Factor Model

Model-implied indicators' covariance matrix of the...

...composite factor model:

$$\Sigma = \begin{pmatrix} \underline{x_1} & \underline{x_2} & \underline{y} & \underline{z} \\ \lambda_1^2 + \theta_1 & \lambda_1\lambda_2 + \theta_{12} & \lambda_1\sigma_{cy} & \lambda_1\sigma_{cz} \\ \lambda_2^2 + \theta_2 & \lambda_2\sigma_{cy} & \lambda_2\sigma_{cz} & \\ \sigma_{yy} & \sigma_{yz} & \sigma_{zz} \end{pmatrix}$$

...common factor model:

$$\Sigma = \begin{pmatrix} \underline{x_1} & \underline{x_2} & \underline{y} & \underline{z} \\ \lambda_1^2 + \theta_1 & \lambda_1\lambda_2 & \lambda_1\sigma_{cy} & \lambda_1\sigma_{cz} \\ \lambda_2^2 + \theta_2 & \lambda_2\sigma_{cy} & \lambda_2\sigma_{cz} & \\ \sigma_{yy} & \sigma_{yz} & \sigma_{zz} \end{pmatrix}$$

⇒ The common factor model is nested in the composite model
[Henseler et al. 2014]

Identification of the Composite Model

Identification of composite models is straightforward:¹

- ▶ Normalization of the weights, e.g., $\mathbf{w}_j' \boldsymbol{\Sigma}_{jj} \mathbf{w}_j = 1$
- ▶ Each composite must be connected to at least one composite or variable not forming the composite

→ All model parameters can be uniquely retrieved from the population indicator covariance matrix

In case of composites embedded in a structural model, also the structural model needs to be identified [Dijkstra, 2017]

¹We ignore trivial regularity assumptions such as weight vectors consisting of zeros only; and similarly, we ignore cases where intra-block covariance matrices are singular.

Model Identification: Degrees of Freedom

For the composite model the degrees of freedom are calculated as follows:

- df = # non-redundant off-diagonal elements of the indicator covariance matrix
- # free correlations among the composites
- # free covariances between the composites and indicators not forming a composite
- # covariances among the indicators not forming a composite
- # free non-redundant off-diagonal elements of each intra-block covariance matrix
- # weights
- + # blocks

For our minimal composite example:

$$df = 6 - 0 - 2 - 1 - 1 - 2 + 1 = 1$$

Estimation of the Composite Model

To determine the weights, several methods have been proposed:

- ▶ Predetermined weights such as unit weights or weights obtained by experts
- ▶ Approaches to generalized canonical correlation analysis (GCCA) such as MAXVAR [Kettenring, 1971]
- ▶ Partial least squares path modeling (PLS-PM) [Wold, 1975]
- ▶ Generalized structured component analysis (GSCA) [Hwang & Takane, 2004]

GCCA: MAXVAR

MAXVAR maximizes the largest eigenvalue of the composite correlation matrix to obtain the weights

Advantage over other approaches to GCCA that it has a closed form expression

Assessment of the Composite Model

The overall model fit can be assessed in two non-exclusive ways:

- ▶ Measures of fit (heuristic rules)
- ▶ Test for overall model fit

Fit Measures

The overall model fit can be assessed in two non-exclusive ways:

- ▶ Standardized root mean squared residual (SRMR)
- ▶ Root mean squared residual covariance matrix (RMS_{Θ})
- ▶ Normed fit index (NFI)
- ▶ ...

More research is required to assess their performance in case of composite models


Test for Overall Model Fit

To test the overall model fit, a bootstrap-based test can be used ($H_0 : \Sigma = \Sigma(\theta)$) [Beran & Srivastava, 1985, Bollen & Stine, 1992] in combination with various discrepancy measures such as


- ▶ SRMR
- ▶ Geodesic distance
- ▶ Squared Euclidean distance

Test for Overall Model Fit

It compares the model-implied indicators' covariance matrix of the composite and a saturated model:



(c) Composite model



(d) Saturated model

If the test is not rejected empirical evidence for the usefulness of the artifact is obtained

Monte Carlo Simulation

Is the test for overall model fit capable to detect misspecifications in the composite model such as

- ▶ Wrongly assigned indicators
- ▶ Correlations between indicators of different blocks that cannot be fully explained by the composites

⇒ Monte Carlo simulation to assess the performance

Simulation setup:

- ▶ 5 population models
- ▶ weights are calculated by MAXVAR
- ▶ 10,000 runs
- ▶ 200 bootstrap runs
- ▶ normally distributed datasets
- ▶ various sample sizes from 50 to 1,450


Monte Carlo Simulation: Population Models

Experimental condition


Population model

Estimated model



1) No misspecification



2) Confounded indicators



3) Unexplained correlation




Monte Carlo Simulation: Population Models

Experimental condition



Population model

Specified model


4) No misspecification



5) Unexplained correlation




Monte Carlo Simulation: Rejection Rates




Significance level: ▲ 10% ■ 5% • 1%

Monte Carlo Simulation: Rejection Rates



Significance level: ▲ 10% ■ 5% ● 1%


Monte Carlo Simulation: Rejection Rates




Significance level: ▲ 10% ■ 5% ● 1%

Extension: Second-order Composites

Emergent variables that are built of other constructs can be modeled and tested such as an emergent variable built of latent variables [Van Riel et al., 2017] or artifacts [Schuberth et al., in progress]



(e) Composite of common factors



(f) Composite of composites

Extension: Multigroup Comparison

It can be assessed whether emergent variables are built the same way across groups (MICOM) [Henseler et al. 2016].

It can be assessed whether the built emergent variable's behavior is the same across groups, i.e., comparing the model-implied indicator variance-covariance matrix across groups using a permutation test [Klesel et al., in press].

Thank you!

Florian Schuberth
email: f.schuberth@utwente.nl
UNIVERSITY OF TWENTE.

References



Beran, R. & Srivastava, M.S. (1985)

Bootstrap tests and confidence regions for functions of a covariance matrix

The Annals of Statistics 13(1) 95 – 115.



Bollen, K. A. & Stine, R. A. (1992)

Bootstrapping goodness-of-fit measures in structural equation models

Sociological Methods & Research 21(2) 205 – 229.



Dijkstra, T. K. (2017)

The perfect match between a model and a mode. In H. Latan & R. Noonan (Eds.)

Partial Least Squares Path Modeling: Basic Concepts, Methodological Issues and Applications, 55 – 80, Springer.

References



Henseler, J., Dijkstra, T. K., Sarstedt, M., Ringle, C. M., Diamantopoulos, A., Straub, D. W., Ketchen, D. J., Hair, J. F., Hult, G. T. M. & Calantone, R. J. (2014)

Common beliefs and reality about PLS: Comments on Rönkkö and Evermann (2013)

Organizational Research Methods 17 182 – 209.



Henseler, J., Ringle, C., & Sarstedt, M. (2016)

Testing measurement invariance of composites using partial least squares

International Marketing Review 33(3) 405 – 431.



Hwang, H. & Takane, Y. (2004)

Generalized structured component analysis

Psychometrika 69(1) 81 – 99.



Kettenring, J.R. (1971)

Canonical analysis of several sets of variables

Biometrika, 58(3), 433 – 451.

References



Klesel, M., Schuberth, F., Henseler, J., & Niehaves, J. (in press)
A test for multigroup comparison in partial least squares path modeling
Internet Research, 1 – 17.



Schuberth, F., Rademaker, M., & Henseler, J. (in progress)
Dealing with hierarchical models containing composites of composites
using PLS path modeling
Working Paper.



Tenenhuis, A. & Tenenhaus, M. (2011)
Regularized generalized canonical correlation analysis
Psychometrika 76(2) 257 – 284.



Van Riel, A. C. R., Henseler, J., Kemény, I., & Sasovova, Z (2017)
Estimating hierarchical constructs using Partial Least Squares: the case of
second order composites of factors
Industrial Management & Data Systems, 117(3) 459 – 477.

References



Wold, A.O.H. (1975)

Path models with latent variables: The NIPALS approach. In H. Blalock, A. Aganbegian, F. Borodkin, R. Boudon, & V. Capecchi (Eds.) *Quantitative Sociology*, 307 - 357, New York Academic Press.